

Neutrino Masses and Mixings in a Universal Seesaw Mass Matrix Model

Yoshio Koide*

Department of Physics, University of Shizuoka
395 Yada, Shizuoka 422, Japan

Abstract

Neutrino masses and mixings are investigated on the basis of a universal seesaw mass matrix model, in which quark (except for top) and charged lepton mass matrices M_f and neutrino mass matrix M_ν are given by $M_f \simeq m_L M_F^{-1} m_R$ and $M_\nu \simeq m_L M_F^{-1} m_L^T$ ($F = N$), respectively. For a simple model which can successfully describe quark masses and mixings, we find that the observed neutrino data (except for the solar neutrino data) are favor to the intermediate mass scales $O(m_R) = 10^{11}$ GeV and $O(M_F) = 10^{13}$ GeV together with $O(m_L) = 10^2$ GeV. In spite of the largesse of $O(m_R)$, the observed top quark mass can be consistently understood from the would-be seesaw mass matrix with these mass scales.

1. Introduction

One of the most exciting problem in the quark and lepton physics is to give a natural understanding of the observed hierarchical mass spectra of quarks and leptons in relation to the electroweak scale Λ_W . In the conventional models, the fundamental fermions acquire their masses through the symmetry breaking of the electroweak symmetry at the energy scale $\mu = \langle \phi_L^0 \rangle \equiv \Lambda_W = 174$ GeV. However, the observed mass values except for the top quark mass m_t are considerably small compared with Λ_W . Especially, the neutrino masses are invisibly small (or exactly zero). In relation to this problem, we know an interesting mechanism, the so-called “seesaw” mechanism.

The mechanism [1] was first proposed in order to answer the question why neutrino masses are so invisibly small: $M_\nu \simeq -mM^{-1}m^T$, where m is a Dirac mass matrix with the order of the conventional quark and charged lepton masses and M is a Majorana mass matrix of the right-handed neutrino ν_{Ri} (i is family-index). If we suppose $m_\nu \leq 10$ eV and $m = O(m_\tau)$, the Majorana mass M must be larger than 10^8 GeV. Usually, in the most models, the order of M is taken as the order of the unification energy scale.

On the other hand, in order to understand why quark masses are so small compared with the electroweak scale Λ_W , the mechanism was applied to the quarks [2]: $M_f \simeq -m_L M_F^{-1} m_R$, where M_F is a mass matrix of hypothetical heavy fermions F_i . If we take $m_L \sim m_W \sim 10^2$ GeV, $m_R \sim 10^3$ GeV and $m_f \sim$ a few GeV, then the heavy fermion mass M_F must be of the order of 10^5 GeV. However, it seems to be not economical that we have two different mass scales M and M_F . Can we build a model with the same mass scale for M and M_F ? The answer is Yes. Since the neutrino and quark mass matrices M_ν and M_f are given by $M_\nu \simeq -m_L M_F^{-1} m_L^T$ ($F = N$) and $M_f \simeq -m_L M_F^{-1} m_R$ ($F = U, D$), respectively, we can understand the smallness of the neutrino masses by assuming $O(m_L)/O(m_R) = O(m_\nu/m_f)$ ($f = u, d$) [3].

By the way, for the seesaw model for quarks, there seems to be a stumbling block: it seems that the observed top quark mass [4] $m_t \sim \Lambda_W$ apparently takes objection to the application of the seesaw mass matrix model to quarks. However, recently, it has been pointed out [5,6] that the seesaw mass matrix model for quarks is rather preferable to understand why only top quark mass m_t is of the order of the electroweak scale Λ_W and why m_t is so singularly enhanced in the third-family in contrast with $m_u \sim m_d$ in the first family. In the framework of the $SU(2)_L \times SU(2)_R \times U(1)_Y$ gauge model, the 6×6 would-be seesaw mass matrix for

the fermion (f, F) is given by

$$M = \begin{pmatrix} 0 & m_L \\ m_R & M_F \end{pmatrix} = m_0 \begin{pmatrix} 0 & Z_L \\ \kappa Z_R & \lambda Y_f \end{pmatrix}, \quad (1.1)$$

where f_i ($i = 1, 2, 3$: family index) are the ordinary quarks ($f = u, d$) and leptons ($f = \nu, e$), and F_i are hypothetical vector-like heavy fermions corresponding to f_i . The fermions f_i and F_i belong to $f_L = (2, 1)$, $f_R = (1, 2)$, $F_L = (1, 1)$ and $F_R = (1, 1)$ of $SU(2)_L \times SU(2)_R$, respectively. The symmetries $SU(2)_L$ and $SU(2)_R$ are broken by the vacuum expectation values of the Higgs scalars ϕ_L and ϕ_R , i.e., the mass matrices m_L and m_R , respectively. For convenience, in (1.1), we have denoted the matrices m_L , m_R and M_F in terms of the matrices Z_L , Z_R and Y_f , which are of the order of one, i.e., $\text{Tr}(Z_L Z_L^\dagger) = O(1)$, $\text{Tr}(Z_R Z_R^\dagger) = O(1)$ and $\text{Tr}(Y_f Y_f^\dagger) = O(1)$. It is well-known that the mass matrix (1.1) leads to the seesaw expression

$$M_f \simeq -m_L M_F^{-1} m_R, \quad (1.2)$$

for the case $|\lambda| \gg |\kappa| \gg 1$ and $\det M_F \neq 0$. On the contrary, for the case $\det M_F = 0$, the mass spectrum is given by [5,6]

$$\begin{aligned} m_1, m_2 &\sim (\kappa/\lambda) m_0, \\ m_3 &\sim m_0 = O(m_L), \\ m_4 &\sim \kappa m_0 = O(m_R), \\ m_5, m_6 &\sim \lambda m_0 = O(M_F), \end{aligned} \quad (1.3)$$

independently of the explicit structures of Z_L , Z_R and Y_f . Note that the third fermion mass m_3^f is given by $m_3^f \sim m_0$ without the suppression factor κ/λ . Therefore, if we build a model such as $\det M_F = 0$ in up-quark sector, we can understand why top-quark alone has a mass of the order of m_L ($\sim \Lambda_W$). This fact (1.3) was first explicitly demonstrated by Fusaoka and the author [5] on the basis of a special seesaw mass matrix model, where M_F is given by the form [(unit matrix)+(a rank-one matrix)], and then re-stressed by Morozumi et al. [6], on the basis of a general study.

Thus, the result (1.3) seems to support the idea that the seesaw mass matrix model for the quarks should be taken seriously. Then, the problem is rather in the neutrino sector: Can the seesaw model, which can successfully describe the quark masses and mixings, satisfactorily describe the neutrino masses and mixings, too? The purpose of the present paper is to discuss the neutrino phenomenology on the

bases of an explicit model which can satisfactorily describe the quark masses and mixings.

In the next section, we discuss the 12×12 mass matrix of the twelve Majorana neutrinos and we will re-derive the well-known form $M_\nu \simeq -(m_0/\lambda)Z_L Y_\nu^{-1} Z_L^T$ for the conventional light neutrinos. In Sec. III, we introduce a model (Model I) with $Z_L = Z_R$ and $Y_\nu = [(\text{unit matrix}) + (\text{rank-one matrix})]$ as a simple example of the matrices Z_L , Z_R and Y_F . In Sec. IV, we try to explain all of the present neutrino data, the solar, atmospheric and LSND neutrino data, by Model I. However, we will fail in the simultaneous explanation of the three data, and we can give a satisfactory explanation only by giving up the explanation of one of the three. For example, if we give up an explanation of the solar neutrino data, we can give the reasonable values of $(\Delta m^2, \sin^2 2\theta)$ for the atmospheric and LSND neutrinos, and we obtain the intermediate mass scales $\kappa m_0 \sim 10^{11}$ GeV and $\lambda m_0 \sim 10^{13}$ GeV together with $m_0 = 312$ GeV. In spite of the largeness of $O(m_R)$, the model will be able to give a reasonable value of the top quark mass m_t (and, of course, reasonable other quark masses and mixings). Finally, Sec. V is devoted to the conclusions and remarks.

2. Neutrino Mass Matrix

The neutral lepton mass matrix which is sandwiched between $(\bar{\nu}_L, \bar{\nu}_R^c, \bar{N}_L, \bar{N}_R^c)$ and $(\nu_L^c, \nu_R, N_L^c, N_R)^T$, where $\nu_L^c \equiv (\nu_L)^c \equiv C\bar{\nu}_L^T$ and so on, is given by

$$M = \begin{pmatrix} 0 & 0 & 0 & m_L \\ 0 & 0 & m_R^T & 0 \\ 0 & m_R & M_M & M_D \\ m_L^T & 0 & M_D^T & M_M \end{pmatrix}, \quad (2.1)$$

where $M_D (\equiv M_N)$ and M_M are Dirac and Majorana mass matrices of the neutral heavy fermions N_i . The matrices m_L and m_R are universal for all fermion sectors, i.e., $f = u, d, e, \nu$, and the differences among up-/down- quark/lepton masses are generated by the differences of $M_F \equiv \lambda Y_f$. The heavy fermions F_i belong to $(1, 1)$ of $SU(2)_L \times U(2)_R$. Besides, the neutral heavy leptons N_i do not have the $U(1)$ -charge. Therefore, it is likely that when the Dirac masses $(M_D)_{ij}$ are generated between $\bar{N}_{Li}$ and N_{Rj} , the Majorana masses $(M_M)_{ij}$ are also generated between $\bar{N}_{Li}$ and N_{Lj}^c ($\bar{N}_{Ri}^c$ and N_{Rj}) with the same structure at the same energy scale $\mu = \lambda m_0$. Hereafter, we put $M_M = M_D \equiv M_N \equiv \lambda m_0 Y_\nu$.

For the case $M_M = M_D$, the diagonalization must be done carefully because the determinant of the 6×6 sub-matrix for (N_L^c, N_R) in the 12×12 matrix (2.1)

becomes zero. First, we rotate the matrix (2.1) on the (N_L^c, N_R) -plane by the angle $\pi/4$. Then, the mass matrix (2.1) becomes

$$M' = \frac{m_0}{\sqrt{2}} \begin{pmatrix} 0 & 0 & -Z_L & Z_L \\ 0 & 0 & \kappa Z_R^T & \kappa Z_R^T \\ -Z_L^T & \kappa Z_R & 0 & 0 \\ Z_L^T & \kappa Z_R & 0 & 2\sqrt{2}\lambda Y_\nu \end{pmatrix}, \quad (2.2)$$

We can see that, of the twelve components of the neutrinos, the three are approximately described by the mass matrix $2\lambda m_0 Y_\nu$. The remaining 9×9 mass matrix is given by

$$M' \simeq \frac{m_0}{\sqrt{2}} \begin{pmatrix} 0 & 0 & -Z_L \\ 0 & 0 & \kappa Z_R^T \\ -Z_L^T & \kappa Z_R & 0 \end{pmatrix} - \frac{m_0}{4\lambda} \begin{pmatrix} Z_L Y^{-1} Z_L^T & \kappa Z_L Y^{-1} Z_R & 0 \\ \kappa Z_R^T Y^{-1} Z_L^T & \kappa^2 Z_R^T Y^{-1} Z_R & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad (2.3)$$

where we have used the formula of the seesaw approximation for the $(n+m) \times (n+m)$ matrix M :

$$M = \begin{pmatrix} A & B \\ C & D \end{pmatrix} \implies M' \simeq \begin{pmatrix} A - BD^{-1}C & 0 \\ 0 & D \end{pmatrix}. \quad (2.4)$$

By using

$$R \simeq \begin{pmatrix} 1 & \varepsilon & 0 \\ -\varepsilon^\dagger & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}, \quad \varepsilon = \frac{1}{\kappa} Z_L (Z_R^T)^{-1}, \quad (2.5)$$

the matrix (2.3) is transformed into

$$RM'R^T \simeq m_0 \begin{pmatrix} -\frac{1}{\lambda} Z_L Y^{-1} Z_L^T & -\frac{\kappa}{2\lambda} Z_L Y^{-1} Z_R & 0 \\ -\frac{\kappa}{2\lambda} Z_R^T Y^{-1} Z_L & -\frac{\kappa^2}{4\lambda} Z_R^T Y^{-1} Z_R & \frac{\kappa}{\sqrt{2}} Z_R \\ 0 & \frac{\kappa}{\sqrt{2}} Z_R & 0 \end{pmatrix}. \quad (2.6)$$

Therefore, we obtain the following twelve Majorana neutrinos: (i) three heavy Majorana neutrinos with masses of the order of λm_0 , whose mass matrix is approximately given by

$$M_{heavy} \simeq 2M_N = 2\lambda m_0 Y_\nu , \quad (2.7)$$

(ii) three sets of almost degenerate two Majorana neutrinos (the pseudo-Dirac neutrino [7]) with masses of the order of κm_0 , whose mass matrix is approximately given by

$$M_{PS-D} \simeq \begin{pmatrix} -\frac{1}{4}m_R^T M_N^{-1} m_R & \frac{1}{\sqrt{2}}m_R^T \\ \frac{1}{\sqrt{2}}m_R & 0 \end{pmatrix} = \kappa m_0 \begin{pmatrix} -\frac{\kappa}{4\lambda} Z_R^T Y_\nu^{-1} Z_R & \frac{\kappa}{\sqrt{2}} Z_R^T \\ \frac{\kappa}{\sqrt{2}} Z_R & 0 \end{pmatrix} , \quad (2.8)$$

and (iii) three light Majorana neutrinos with masses of the order of $(1/\lambda)m_0$, whose mass matrix is approximately given by

$$M_\nu \simeq -m_L M_N^{-1} m_L^T = -\frac{m_0}{\lambda} Z_L Y_\nu^{-1} Z_L^T . \quad (2.9)$$

Note that, differently from the models by Berezhiani and Davidson-Wari [3], there are no neutrinos with masses of the order of $(\kappa^2/\lambda)m_0$, i.e., whose mass matrix is approximately given by $m_R M_N^{-1} m_R^T$.

The neutrinos which are described by the mass matrix (2.9) consist of almost left-handed neutrinos ν_{Li} . Therefore, our task is to seek for such matrix forms of Z_L , Z_R and Y_f as those can give reasonable quark and lepton masses, Cabibbo-Kobayashi-Maskawa (CKM) [8] matrix, and neutrino mixings, where the mass matrices of down-quarks and charged leptons are given by (1.2), i.e., $M_f \simeq (\kappa/\lambda) Z_L Y_f^{-1} Z_R$ with $\det Y_f \neq 0$, that of up-quarks is given by (1.1) with $\det Y_u = 0$, and that of neutrinos is given by (2.9).

3. A Simple Case

As an explicit model of the universal seesaw mass matrix models which give phenomenologically reasonable predictions for quarks, there is a model which has been proposed by Fusaoka and the author [5]. In their model, the matrices are simply taken by

$$Z_L = Z_R \equiv Z , \quad (3.1)$$

and

$$Y_f = \mathbf{1} + 3b_f X , \quad (3.2)$$

where $\mathbf{1}$ and X are a 3×3 unit matrix and a rank-one matrix normalized as $X^2 = X$, respectively. They have assumed that the matrix Z is given by a diagonal form in the family-basis on which X is democratic form:

$$Z = \begin{pmatrix} z_1 & 0 & 0 \\ 0 & z_2 & 0 \\ 0 & 0 & z_3 \end{pmatrix}, \quad X = \frac{1}{3} \begin{pmatrix} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{pmatrix}. \quad (3.3)$$

They have used the input values

$$\frac{z_1}{\sqrt{m_e}} = \frac{z_2}{\sqrt{m_\mu}} = \frac{z_3}{\sqrt{m_\tau}} = \frac{1}{\sqrt{m_e + m_\mu + m_\tau}}, \quad (3.4)$$

from $M_e \simeq (\kappa/\lambda)m_0 Z \cdot \mathbf{1} \cdot Z$ by assuming $b_e = 0$ in the charged lepton sector as a trial. Then, for up-quark sector with $\det Y_u = 0$, i.e., with $b_u = -1/3$, they have obtained the successful relations

$$m_u \simeq \frac{3}{2} \frac{m_e}{m_\tau} \frac{\kappa}{\lambda} m_0, \quad m_c \simeq 2 \frac{m_\mu}{m_\tau} \frac{\kappa}{\lambda} m_0, \quad m_t \simeq \frac{1}{\sqrt{3}} m_0. \quad (3.5)$$

As we noted in (1.3), the third quark mass does not have the suppression factor κ/λ . The ratio of κ/λ is fixed as $\kappa/\lambda \simeq 0.02$ from the observed ratio of m_c/m_t . For down quark sectors, by choosing the complex parameter b_f as $b_d \simeq -e^{i\pi/10}$, they have obtained reasonable results of quark masses (not only m_i^d/m_j^d , but also m_i^u/m_j^d) and CKM matrix parameters. Hereafter, we will refer this model as Model I.

Therefore, our next interest is whether Model I is applicable to neutrino phenomenology under the same parameter values or not. Although they have taken $\kappa = 10$ tentatively, the parameter κ is essentially free because we does not yet observe the right-handed weak boson. [The case with a small κ (e.g., ~ 10) is very attractive from the phenomenological standpoint, because the case can bring us detectable new physics in abundance [9]. However, in the present paper, since we intend to give a unified description of quark and neutrino mass matrices without assuming further additional intermediate mass scale, we do not consider the case with $\kappa \sim 10$.] Also, although the parameters b_f have been taken as $b_e = 0$, $b_u = -1/3$ and $b_d \simeq -e^{i\pi/10}$ in Ref. [5], the value of b_ν in the neutrino sector is still free. Therefore, for the neutrino masses and mixings, we have three adjustable parameters, i.e., b_ν (complex) and m_0/λ (real).

For typical values of b_ν , $b_\nu \simeq -1/3$, $b_\nu \simeq -1/2$ and $b_\nu \simeq -1$, the masses m_i^ν ($i = 1, 2, 3$) and mixings $U_{\alpha i}$ ($\alpha = e, \mu, \tau$; $i = 1, 2, 3$) for three light neutrinos are given as follows [10].

Case $b_\nu = -1/3 + \Delta b_\nu$ ($1 \gg |\Delta b_\nu| \neq 0$):

$$m_1^\nu \simeq \frac{3}{2} \frac{m_e}{m_\tau} \frac{m_0}{\lambda}, \quad m_2^\nu \simeq 2 \frac{m_\mu}{m_\tau} \frac{m_0}{\lambda}, \quad m_3^\nu \simeq \frac{2\sqrt{2}}{27|\Delta b_\nu|} \frac{m_0}{\lambda}, \quad (3.6)$$

$$U \simeq \begin{pmatrix} 1 & \frac{1}{2}\sqrt{\frac{m_e}{m_\mu}} & \sqrt{\frac{m_e}{m_\tau}} \\ -\frac{1}{2}\sqrt{\frac{m_e}{m_\mu}} & 1 & \sqrt{\frac{m_\mu}{m_\tau}} \\ -\frac{1}{2}\sqrt{\frac{m_e}{m_\tau}} & -\sqrt{\frac{m_\mu}{m_\tau}} & 1 \end{pmatrix}, \quad (3.7)$$

Case $b_\nu \simeq -1/2$:

$$m_1^\nu \simeq 2 \frac{m_e}{m_\tau} \frac{m_0}{\lambda}, \quad m_2^\nu \simeq m_3^\nu \simeq \sqrt{\frac{m_\mu}{m_\tau}} \frac{m_0}{\lambda}, \quad (3.8)$$

$$U \simeq \begin{pmatrix} 1 & \sqrt{\frac{m_e}{2m_\mu}} & \sqrt{\frac{m_e}{2m_\mu}} \\ -\sqrt{\frac{m_e}{m_\mu}} & \frac{1}{\sqrt{2}} & \mp \frac{1}{\sqrt{2}} \\ -\sqrt{\frac{m_e}{m_\tau}} & \pm \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \end{pmatrix}, \quad (3.9)$$

Case $b_\nu \simeq -1$:

$$m_1^\nu \simeq m_2^\nu \simeq \sqrt{\frac{m_e m_\mu}{m_\tau^2}} \frac{m_0}{\lambda}, \quad m_3^\nu \simeq \frac{1}{2} \frac{m_0}{\lambda}, \quad (3.10)$$

$$U \simeq \begin{pmatrix} \frac{1}{\sqrt{2}} & \mp \frac{1}{\sqrt{2}} & -\sqrt{\frac{m_e}{m_\tau}} \\ \pm \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} & -\sqrt{\frac{m_\mu}{m_\tau}} \\ \sqrt{\frac{m_\mu}{2m_\tau}} & \sqrt{\frac{m_\mu}{2m_\tau}} & 1 \end{pmatrix} . \quad (3.11)$$

4. Neutrino Data and their Interpretations

As possible evidences for non-zero neutrino masses, at present, the following data are known:

(a) The solar neutrino data [11] with the Mikheyev-Smirnov-Wolfenstein (MSW) effect [12] have suggested

$$\Delta m_\odot^2 \simeq 6 \times 10^{-6} \text{ eV}^2 , \quad \sin^2 2\theta_\odot \simeq 7 \times 10^{-3} , \quad (4.1)$$

(the small-angle solution), or

$$\Delta m_\odot^2 \simeq 10^{-5} \text{ eV}^2 , \quad \sin^2 2\theta_\odot \simeq 0.8 , \quad (4.2)$$

(the large-angle solution).

(b) The atmospheric neutrino data reported by the Kamiokande collaboration [13] have suggested a neutrino mixing $\nu_\mu \leftrightarrow \nu_x$:

$$\Delta m_{atm}^2 \simeq 1.6 \text{ (1.8)} \times 10^{-2} \text{ eV}^2 , \quad \sin^2 2\theta_{atm} \simeq 1 , \quad (4.3)$$

for $x = \mu$ ($x = e$).

(c) The neutrino oscillation ($\bar{\nu}_\mu \rightarrow \bar{\nu}_e$) experiment by the liquid scintillator neutrino detector (LSND) [14] at Los Alamos has been reported nonzero neutrino mass:

$$(\Delta m^2, \sin^2 2\theta)_{LSND} \simeq (0.3 \text{ eV}^2, 0.04) - (2 \text{ eV}^2, 0.002) . \quad (4.4)$$

(d) A cosmological model with cold+hot dark matter (CHDM) suggests [15]

$$m_1^\nu + m_2^\nu + m_3^\nu \simeq 4.8 \text{ eV} . \quad (4.5)$$

These mass values and mixings (4.1)–(4.3) and (4.5) are not based on direct observations of masses and mixings and they are highly model-dependent. On the other hand, the experiment which reported the result (4.4) is still controversial [16]. (For

convenience, hereafter, we will refer these data (4.1)–(4.5) as $\nu_{\odot}^{small}$, $\nu_{\odot}^{large}$, ν_{atm} , ν_{LSND} and ν_{CHDM} , respectively.)

Since $\Delta m_{LSND}^2 \gg \Delta m_{atm}^2 \gg \Delta m_{\odot}^2$, we investigate only the following two cases: (A) $\Delta m_{32}^2 \gg \Delta m_{21}^2$ (i.e., $b_{\nu} \simeq -1$ and $b_{\nu} \simeq -1/3$); (B) $\Delta m_{32}^2 \ll \Delta m_{21}^2$ (i.e., $b_{\nu} \simeq -1/2$), where $\Delta m_{ij}^2 = m_i^2 - m_j^2$. Of course, in the three family model, we cannot assign the three values Δm_{LSND}^2 , Δm_{atm}^2 and $\Delta m_{\odot}^2$ to $(\Delta m_{32}^2, \Delta m_{21}^2)$ simultaneously. For the case (A), we consider two cases: (A₁) $(\Delta m_{32}^2, \Delta m_{21}^2) = (\Delta m_{LSND}^2, \Delta m_{\odot}^2)$ and (A₂) $(\Delta m_{32}^2, \Delta m_{21}^2) = (\Delta m_{LSND}^2, \Delta m_{atm}^2)$. Similarly, for the case (B), we consider two cases: (B₁) $(\Delta m_{32}^2, \Delta m_{21}^2) = (\Delta m_{\odot}^2, \Delta m_{LSND}^2)$ and $(\Delta m_{32}^2, \Delta m_{21}^2) = (\Delta m_{atm}^2, \Delta m_{LSND}^2)$. We do not consider the cases $(\Delta m_{atm}^2, \Delta m_{\odot}^2)$ and $(\Delta m_{\odot}^2, \Delta m_{atm}^2)$, because it is readily known that the cases cannot give the observed large mixing $\sin^2 2\theta_{atm} \sim 1$ in ν_{atm} .

In the three-family model, the neutrino oscillation $P(\nu_{\alpha} \rightarrow \nu_{\beta})$ ($\alpha \neq \beta$) is given by

$$P(\nu_{\alpha} \rightarrow \nu_{\beta}) = S_{21}^{\alpha\beta} S_{21} + S_{31}^{\alpha\beta} S_{31} + S_{32}^{\alpha\beta} S_{32} \quad , \quad (4.6)$$

where

$$S_{ij}^{\alpha\beta} = -4\Re(U_{\alpha i} U_{\beta i}^* U_{\alpha j}^* U_{\beta j}) \quad , \quad (4.7)$$

$$S_{ij} = \sin^2(L\Delta m_{ij}^2/4E_{\nu}) \quad , \quad (4.8)$$

and we have neglected CP -violation terms in (4.6).

In the case (A₁), since $\Delta m_{32}^2 \gg \Delta m_{atm}^2 \gg \Delta m_{21}^2$, we can regard S_{ij} as $S_{21} \simeq 0$ and $\langle S_{31} \rangle = \langle S_{32} \rangle = 1/2$ for the atmospheric neutrinos, where $\langle S_{ij} \rangle$ denotes the mean value of $S_{ij}(L/E_{\nu})$. Then, by using $P(\nu_{\alpha} \rightarrow \nu_{\beta}) \simeq 2|U_{\alpha 3}|^2|U_{\beta 3}|^2$, the ratio

$$R_{atm} = \frac{(\nu_{\mu}/\nu_e)_{data}}{(\nu_{\mu}/\nu_e)_{MC}} \simeq \frac{P(\nu_{\mu} \rightarrow \nu_{\mu}) + \frac{1}{2}P(\nu_e \rightarrow \nu_{\mu})}{P(\nu_e \rightarrow \nu_e) + 2P(\nu_{\mu} \rightarrow \nu_e)} \quad , \quad (4.9)$$

is expressed as

$$R_{atm} \simeq \frac{1 - |U_{e3}|^2|U_{\mu 3}|^2 - 2|U_{\mu 3}|^2|U_{\tau 3}|^2}{1 + 2|U_{e3}|^2|U_{\mu 3}|^2 - 2|U_{e3}|^2|U_{\tau 3}|^2} \quad . \quad (4.10)$$

Therefore, in the case (A₁), instead of (4.3), we temporize with (4.10) whose value is consistent with the observed value [13]

$$R_{atm} = 0.57_{-0.07}^{+0.08} \pm 0.07 \quad . \quad (4.11)$$

However, both cases $b_{\nu} \simeq -1/3$ and $b_{\nu} \simeq -1$, from (3.7) and (3.11), respectively, give $R_{atm} \simeq 1 - 2m_{\mu}/m_{\tau} \simeq 1$, which is in disagreement with the observed value

(4.11). On the other hand, the mixing parameter $\sin^2 2\theta_\odot$ in the MSW solutions is interpreted by

$$\sin^2 2\theta_\odot = 4|U_{e1}|^2|U_{e2}|^2 . \quad (4.12)$$

The cases $b_\nu \simeq -1/3$ and $b_\nu \simeq -1$ give

$$\sin^2 2\theta_\odot \simeq m_e/m_\mu \simeq 0.005 , \quad (4.13)$$

and

$$\sin^2 2\theta_\odot \simeq 1 , \quad (4.14)$$

which are consistent with the small- and large-angle solutions (4.1) and (4.2), respectively.

In the case (A₂), since $\Delta m_\odot^2 \ll \Delta m_{21}^2 \ll \Delta m_{32}^2$, we can regard S_{ij} as $\langle S_{21} \rangle = \langle S_{31} \rangle = \langle S_{32} \rangle = 1/2$ for $\nu_\odot$. Then, we obtain

$$P_{ee} \equiv P(\nu_e \rightarrow \nu_e) = |U_{e1}|^4 + |U_{e2}|^4 + |U_{e3}|^4 . \quad (4.15)$$

According to the recent analysis by Acker and Pakvasa [17] we will search for a solution which is consistent with

$$P_{ee} = 0.42 - 0.52 . \quad (4.16)$$

The cases $b_\nu \simeq -1/3$ and $b_\nu \simeq -1$, from (3.7) and (3.11), give $P_{ee} \simeq 1$ and $P_{ee} \simeq 1/2$, respectively. Therefore, only the case $b_\nu \simeq -1$ can explain the solar neutrino data. On the other hand, the mixing parameter $\sin^2 2\theta_{atm}$ is interpreted by

$$\sin^2 2\theta_{atm} = S_{21}^{\mu e} = -4\Re(U_{\mu 2}U_{e2}^*U_{\mu 1}^*U_{e1}) . \quad (4.17)$$

The cases $b_\nu \simeq -1/3$ and $b_\nu \simeq -1$ give $\sin^2 2\theta_{atm} \simeq m_e/m_\mu \simeq 0.005$ and $\sin^2 2\theta_{atm} \simeq 1$, respectively. Again, only the case $b_\nu \simeq -1$ is favorable to the data.

In both cases (A₁) and (A₂), the mixing parameter $\sin^2 2\theta_{LSND}$ is given by

$$\sin^2 2\theta_{LSND} = 4|U_{e3}|^2|U_{\mu 3}|^2 . \quad (4.18)$$

The cases $b_\nu \simeq -1/3$ and $b_\nu \simeq -1$ give $\sin^2 2\theta_{LSND} \simeq 4m_em_\mu/m_\tau^2 \simeq 7 \times 10^{-5}$, which is too small compared with the observed mixing value (4.4).

Similarly, for the case (B₁), we use the expressions

$$\sin^2 2\theta_\odot = 4|U_{e3}|^2|U_{e2}|^2 . \quad (4.19)$$

$$R_{atm} \simeq \frac{1 - |U_{e1}|^2|U_{\mu 1}|^2 - 2|U_{\mu 1}|^2|U_{e1}|^2}{1 + 2|U_{e1}|^2|U_{\mu 1}|^2 - 2|U_{e1}|^2|U_{\tau 1}|^2} . \quad (4.20)$$

The mixing matrix (3.9) gives $\sin^2 2\theta_\odot \simeq (m_e/m_\mu)^2 \simeq 2 \times 10^{-5}$ and $R_{atm} \simeq 1 - m_e/2m_\mu \simeq 1$, which are in disagreement with the data.

In the case (B₂), the solar neutrino data is explained by P_{ee} given by (4.15), and the mixing parameter in ν_{atm} is expressed by

$$\sin^2 2\theta_{atm} = S_{23}^{\mu\tau} = -4Re(U_{\mu 2}U_{\tau 2}^*U_{\mu 3}^*U_{\tau 3}) . \quad (4.21)$$

Since the mixing matrix (3.9) gives $P_{ee} \simeq 1$ and $\sin^2 2\theta_{atm} \simeq 1$, we fail to explain the $\nu_\odot$ data, but we can understand the large mixing in ν_{atm} .

For both cases (B₁) and (B₂), the mixing parameter $\sin^2 2\theta_{LSND}$ is given by

$$\sin^2 2\theta_{LSND} = 4|U_{e1}|^2|U_{\mu 1}|^2 \simeq 4m_e/m_\mu \simeq 0.02 , \quad (4.22)$$

where we have used the mixing matrix (3.9) for $b_\nu \simeq -1/2$. The mixing value (4.22) corresponds to $\Delta m_{LSND}^2 \simeq 0.5 \text{ eV}^2$ in the LSND allowed region (4.4).

The results are summarized in Table I. We find that the present model cannot give a simultaneous interpretation for the three neutrino data, $\nu_\odot$, ν_{atm} and ν_{LSND} . We must give up the explanation of one of the three data.

As suggested by the case (B₂) in Table I, if we give up explaining the solar neutrino data, we can find an interesting solution: $(\Delta m_{32}^2, \Delta m_{21}^2) = (\Delta m_{atm}^2, \Delta m_{LSND}^2)$ with $b_\nu \simeq -1/2$. For example, the solution $b_\nu = -(1/2)e^{i\beta_\nu}$ with $b_\nu = 0.12^\circ$ gives

$$\sin^2 2\theta_{atm} = 0.995 , \quad \sin^2 2\theta_{LSND} = 0.0191 , \quad (4.23)$$

$$\begin{aligned} m_1^\nu &= 0.000540 \text{ } m_0/\lambda , \\ m_2^\nu &= 0.2288 \text{ } m_0/\lambda , \\ m_3^\nu &= 0.2326 \text{ } m_0/\lambda , \end{aligned} \quad (4.24)$$

$$r \equiv \Delta m_{32}^2/\Delta m_{21}^2 = 0.0331 . \quad (4.25)$$

The input value $\Delta m_{32}^2 = \Delta m_{atm}^2 = 0.016 \text{ eV}^2$ predicts

$$m_1^\nu = 0.0016 \text{ eV} , \quad m_2^\nu = 0.695 \text{ eV} , \quad m_3^\nu = 0.707 \text{ eV} , \quad (4.26)$$

$$\Delta m_{21}^2 = 0.483 \text{ eV}^2 , \quad (4.27)$$

together with

$$m_0/\lambda = 3.04 \text{ eV} . \quad (4.28)$$

The values (4.23) and (4.27) are in good agreement with the observed LSND solution $(\Delta m^2, \sin^2 2\theta)_{LSND} \simeq (0.5 \text{ eV}^2, 0.02)$. Although the sum $\sum m_i^\nu = 1.4 \text{ eV}$ is

somewhat small compared with the mass value in the CHDM scenario (4.5), these neutrinos can still be one of the dark matter objects

When the LSND data are neglected, the case (A₂) with $b_\nu \simeq -1$ is also interesting. For example, the parameter value $b_\nu = -e^{i\beta_\nu}$ with $\beta_\nu = 1.8^\circ$ gives

$$P_{ee} = 0.52 , \quad \sin^2 2\theta_{atm} = 0.90 , \quad (4.29)$$

which are favorable to the data of $\nu_\odot$ and ν_{atm} . On the other hand, the input

$$\Delta m_{21}^2 = 1.283 \times 10^{-5} (m_0/\lambda)^2 = \Delta m_{atm}^2 = 1.8 \times 10^{-2} \text{ eV}^2 , \quad (4.30)$$

leads to

$$\begin{aligned} m_1^\nu &= 0.00305 \, m_0/\lambda = 0.114 \text{ eV} , \\ m_2^\nu &= 0.00471 \, m_0/\lambda = 0.176 \text{ eV} , \\ m_3^\nu &= 0.5002 \, m_0/\lambda = 18.7 \text{ eV} , \end{aligned} \quad (4.31)$$

together with

$$m_0/\lambda = 37.5 \text{ eV} . \quad (4.32)$$

The value $\sum m_i^\nu \simeq 19 \text{ eV}$ is somewhat large to identify these neutrinos as those in the dark matter scenario. In the present case, although the third neutrino has a considerably large value $m_3^\nu = 18.7 \text{ eV}$, the effective electron neutrino mass $\langle m_\nu \rangle$ is safely small compared with the upper bound on $\langle m_\nu \rangle$ from the neutrinoless double beta decays, $\langle m_\nu \rangle < 0.68 \text{ eV}$ [18], because of the smallness of $|U_{e3}^2|$, i.e.,

$$\langle m_\nu \rangle = |m_1^\nu U_{e1}^2 + m_2^\nu U_{e2}^2 + m_3^\nu U_{e3}^2| = 0.0027 \text{ eV} . \quad (4.33)$$

In these examples, we have chosen the solutions with $\beta_\nu \neq 0$. However, this is not essential for the numerical results. The similar results to the cases $b_\nu = -(1/2) \exp[i 0.12^\circ]$ and $b_\nu = -\exp[i 1.8^\circ]$ can also be obtained by choosing $b_\nu = -0.50004$ and $b_\nu = -0.97$. However, since we have taken the rational solutions of $|b_f|$ in the quark and charged lepton sectors, i.e., $|b_u| = 1/3$, $|b_d| = 1$ and $|b_e| = 0$ [5], we have taken the rational solutions of $|b_\nu|$ in Table II.

In Table II, we summarize these typical cases. For reference, in Table II, we show the values of $\sin^2 2\theta^{\mu\tau}$ which is given by $\sin^2 2\theta^{\mu\tau} = 4|U_{\mu 3}|^2|U_{\tau 3}|^2$ for the case $\Delta m_{32}^2 \gg \Delta m_{21}^2$ and by $\sin^2 2\theta^{\mu\tau} = 4|U_{\mu 1}|^2|U_{\tau 1}|^2$ for the case $\Delta m_{32}^2 \ll \Delta m_{21}^2$, because CHORUS [19] and NOMAD [20] experiments at CERN are expected to present results on the $\nu_\mu \rightarrow \nu_\tau$ oscillation in the very near future. As pointed out by

Tanimoto [21], the values of $\sin^2 2\theta^{\mu\tau}$ are very small for the case $m_1^\nu \ll m_2^\nu \simeq m_3^\nu$, i.e., $\sin^2 2\theta^{\mu\tau} \simeq 4m_e/m_\tau$ for $b_\nu \simeq -1/2$, so that it will be hopeless to observe the $\nu_\mu \rightarrow \nu_\tau$ oscillation at CHORUS and NOMAD. The case $\Delta m_{32}^2 \gg \Delta m_{21}^2$, i.e., $b_\nu \simeq -1$, gives a sizable value $\sin^2 2\theta^{\mu\tau} \simeq 4m_\mu/m_\tau$, so that the case will be hopeful to observe the evidence of the $\nu_\mu \rightarrow \nu_\tau$ oscillation.

In Table II, we also list the value of the intermediate mass scales κm_0 and λm_0 , which are obtained by the following relations:

$$m_t \simeq \frac{1}{\sqrt{3}} m_0 \simeq 180 \text{ GeV} \quad (\text{at } \mu = m_Z) , \quad (4.34)$$

from (3.5) and

$$(\kappa/\lambda)m_0 = m_\tau + m_\mu + m_e = 1.850 \text{ GeV} , \quad (\text{at } \mu = m_Z) \quad (4.35)$$

from $M_e \simeq (\kappa/\lambda)m_0 Z \cdot \mathbf{1} \cdot Z$, i.e.,

$$m_0 \simeq \sqrt{3} m_t \simeq 312 \text{ GeV} \quad (\text{at } \mu = m_Z) , \quad (4.36)$$

$$\kappa/\lambda = 5.93 \times 10^{-3} . \quad (4.37)$$

5. Conclusions

In conclusion, we have investigated a universal seesaw mass matrix model with a form (1.1), and have found that the model has a possibility of giving a unified description of a mass hierarchy of quarks and leptons by assuming $\det M_F = 0$ for the heavy fermion mass matrix M_F in the up-quark sector. Here, the ‘‘hierarchy’’ discussed in the present paper means the mass hierarchy among the three groups, (i) top quark, (ii) quarks except for top and charged leptons, and (iii) neutrinos. For example, in Model I, the hierarchy among m_e , m_μ and m_τ are given by hand, i.e., by adjusting the parameters z_i in the order-one matrices $Z \equiv Z_L = Z_R$. It is our task at the next step to answer why these matrices Z_L , Z_R and Y_f take such structures.

In the $SU(2)_L \times SU(2)_R \times U(1)_Y$ model, it is likely that the neutral heavy leptons acquire Majorana masses M_M together with the Dirac masses M_N with the same structure and the same magnitude, i.e., $M_M = M_N$. Then, masses of the twelve Majorana neutrinos are given by (2.7)–(2.9). In the present model, heavy neutrinos with masses of the order of $(\kappa^2/\lambda)m_0$ do not appear, and, instead, pseudo-Dirac neutrinos with masses of the order of κm_0 appear.

A suitable choice of the matrices Z_L , Z_R and Y_f will give a unified descriptions of masses and mixings of quarks and leptons. As a simple example of Z_L , Z_R and Y_f , a model with (3.1)–(3.3), Model I, has been investigated. The model can give a unified description of masses and mixings of quarks and charged leptons [5]. However, the straightforward application to the neutrino phenomenology fails to give the simultaneous explanation of the solar, atmospheric and LSND neutrino data. If we give up explaining one of these three data, for example, if we give up the explanation of the solar neutrino data, we obtain the intermediate mass scales $\kappa m_0 \sim 10^{11}$ GeV and $\lambda m_0 \sim 10^{13}$ GeV together with $m_0 = 312$ GeV. Note that in spite of the largeness of $O(m_R)$, the model can give a reasonable value of the top quark mass m_t . (In other words, the above intermediate mass scales have estimated from the input value $m_t(m_Z) = 180$ GeV.)

If we take all of the neutrino data, $\nu_\odot$, ν_{atm} and ν_{LSND} seriously, we must seek another set of the matrices Z_L , Z_R and Y_f or we must abandon the idea $O(M_N) = O(M_F)$ ($F = E, U, D$). Although the model with $Z_L = Z_R$, Model I, has successfully described the quark masses and their mixings in terms of charged lepton masses, there is no reason that we should consider $Z_L = Z_R$. It will be worth while investigating a model with $Z_L \neq Z_R$ for quarks and leptons. It will also be worth while investigating a model with $O(M_N) \gg O(M_F)$ ($F = E, U, D$). The latter possibility will bring abundant light neutrinos into the model (for example, see a model given in [10]), although it is not economical because it bring additional intermediate mass scale into the model.

Model I discussed in Secs. III and IV is only an example. There will be many interesting versions of the universal seesaw model. The universal seesaw model will be one of the most promising models of the quark and lepton unification.

Acknowledgments

The author would like to thank M. Tanimoto for his helpful comments and calling his attention to CHORUS and NOMAD experiments. He also thank N. Muto for numerical check of the approximate expressions (2.7)–(2.9) of the 12×12 would-be seesaw mass matrix (2.1) by his computer program with high accuracy. This work was supported by the Grant-in-Aid for Scientific Research, the Ministry of Education, Science and Culture, Japan (No.08640386).

References

- [1] M. Gell-Mann, P. Rammond and R. Slansky, in *Supergravity*, edited by P. van Nieuwenhuizen and D. Z. Freedman (North-Holland, 1979); T. Yanagida, in

Proc. Workshop of the Unified Theory and Baryon Number in the Universe, edited by A. Sawada and A. Sugamoto (KEK, 1979); R. Mohapatra and G. Senjanovic, Phys. Rev. Lett. **44**, 912 (1980).

- [2] Z. G. Berezhiani, Phys. Lett. **129B**, 99 (1983); Phys. Lett. **150B**, 177 (1985); D. Chang and R. N. Mohapatra, Phys. Rev. Lett. **58**, 1600 (1987); A. Davidson and K. C. Wali, Phys. Rev. Lett. **59**, 393 (1987); S. Rajpoot, Mod. Phys. Lett. **A2**, 307 (1987); Phys. Lett. **191B**, 122 (1987); Phys. Rev. **D36**, 1479 (1987); K. B. Babu and R. N. Mohapatra, Phys. Rev. Lett. **62**, 1079 (1989); Phys. Rev. **D41**, 1286 (1990); S. Ranfone, Phys. Rev. **D42**, 3819 (1990); A. Davidson, S. Ranfone and K. C. Wali, Phys. Rev. **D41**, 208 (1990); I. Sogami and T. Shinohara, Prog. Theor. Phys. **66**, 1031 (1991); Phys. Rev. **D47**, 2905 (1993); Z. G. Berezhiani and R. Rattazzi, Phys. Lett. **B279**, 124 (1992); P. Cho, Phys. Rev. **D48**, 5331 (1994); A. Davidson, L. Michel, M. L. Sage and K. C. Wali, Phys. Rev. **D49**, 1378 (1994); W. A. Ponce, A. Zepeda and R. G. Lozano, Phys. Rev. **D49**, 4954 (1994).
- [3] Z. G. Berezhiani, in Ref. [2]; A. Davidson and K. C. Wali, in Ref. [2]; S. Rajpoot, in Ref. [2]; A. Davidson, S. Ranfone and K. C. Wali, in Ref. [2]; W. A. Ponce, A. Zepeda and R. G. Lozano, in Ref. [2].
- [4] CDF collaboration, F. Abe *et al.*, Phys. Rev. Lett. **73**, 225 (1994).
- [5] Y. Koide and H. Fusaoka, Z. Phys. **C71**, 459 (1966); Prog. Theor. Phys. **97**, 459 (1997).
- [6] T. Morozumi, T. Satou, M. N. Rebelo and M. Tanimoto, Preprint HUPD-9704 (1977), hep-ph/9703249.
- [7] L. Wolfenstein, Nucl. Phys. **B185**, 147 (1981); S. T. Petcov, Phys. Lett. **110B**, 245 (1982); C. N. Leung and S. T. Petcov, Nucl. Phys. **125B**, 461 (1983); M. Doi, M. Kenmoku, T. Kotani, H. Nishiura and E. Takasugi, Prog. Theor. Phys. **70**, 1331 (1983); J. W. F. Valle, Phys. Rev. **D27**, 1672 (1983); J. W. F. Valle and M. Singer, Phys. Rev. **D28**, 540 (1983); D. Wyler and L. Wolfenstein, Nucl. Phys. **B218**, 205 (1983).
- [8] N. Cabibbo, Phys. Rev. Lett. **10**, 531 (1966); M. Kobayashi and T. Maskawa, Prog. Theor. Phys. **49**, 652 (1973).
- [9] Y. Koide, Preprint US-96-09 (1996), hep-ph/9701261 (unpublished); US-97-04 (1997), hep-ph/9706277, to be published in Phys. Rev. **D56**, (1997).

- [10] Y. Koide, Mod. Phys. Lett. **A36**, 2849 (1996).
- [11] GALLEX collaboration, P. Anselmann *et al*, Phys. Lett. **B327**, 377 (1994); **B357**, 237 (1995); SAGE collaboration, J. N. Abdurashitov *et al*, Phys. Lett. **B328**, 234 (1994). Also see, N. Hata and P. Langacker, Phys. Rev. **D50**, 632 (1994); **D52**, 420 (1995).
- [12] S. P. Mikheyev and A. Yu. Smirnov, Yad. Fiz. **42**, 1441 (1985); [Sov. J. Nucl. Phys. **42**, 913 (1985)]; Prog. Part. Nucl. Phys. **23**, 41 (1989); L. Wolfenstein, Phys. Rev. **D17**, 2369 (1978); **D20**, 2634 (1979); T. K. Kuo and J. Pantaleon, Rev. Mod. Phys. **61**, 937 (1989). Also see, A. Yu. Smirnov, D. N. Spergel and J. N. Bahcall, Phys. Rev. **D49**, 1389 (1994).
- [13] Y. Fukuda *et al*., Phys. Lett. **B335**, 237 (1994). Also see, Soudan-2 collaboration, M. Goodman *et al*., Nucl. Phys. (Proc. Suppl.) **B38**, 337 (1995); IMB collaboration, D. Casper *et al*, Phys. Rev. Lett. **66**, 2561 (1989); R. Becker-Szendy *et al*, Phys. Rev. **D46**, 3720 (1989).
- [14] C. Athanassopoulos *et al*., Phys. Rev. Lett. **75**, 2650 (1995); Phys. Rev. Lett. **77**, 3082 (1996); nucl-ex/9706006 (1997).
- [15] J. R. Primack, J. Holtzman, A. Klypin and D. O. Caldwell, Phys. Rev. Lett. **74**, 2160 (1995); D. Pogosyan and A. Starobinsky, astro-ph/9502019.
- [16] J. E. Hill, Phys. Rev. Lett. **75**, 2654 (1995).
- [17] A. Acker and S. Pakvasa, Phys. Lett. **B397**, 209 (1997).
- [18] M. K. Moe, Nucl. Phys. (Proc. Suppl.) **B38**, 36 (1995); A. Balysh *et al*., Proc. of the Inst. Conference on High Energy Physics (ICHEP), 27th, Glasgow, Scotland, Eds, P. J. Bussey and I. G. Knowles, IOP, 1995, p.939 (hep-ex/9502007).
- [19] M. Baldo-Ceolin, Nucl. Phys. (Proc. Suppl.) **35**, 450 (1994); K. Winter, *ibid*. **38**, 211 (1995).
- [20] L. DiLella, Nucl. Phys. (Proc. Suppl.) **31**, 319 (1993).
- [21] M. Tanimoto, Phys. Rev. **D53**, 6632 (1996).

Table I. Possible interpretations for the $\nu_\odot$, ν_{atm} and ν_{LSND} data. The parameter r denotes $r = \Delta m_{32}^2 / \Delta m_{21}^2$. The numerical results have obtained from the direct evaluation of the mass matrix (2.1) without using the approximate expressions (3.7), (3.9) and (3.11). The values with the symbols $\circ$ and $\bullet$ denote that those are favorable and unfavorable to the observed data, respectively.

(A) $\Delta m_{32}^2 \gg \Delta m_{21}^2$		(B) $\Delta m_{32}^2 \ll \Delta m_{21}^2$
$b_\nu \simeq -1$	$b_\nu \simeq -1/3$	$b_\nu \simeq -1/2$
(A ₁) $(\Delta m_{32}^2, \Delta m_{21}^2) = (\Delta m_{LSND}^2, \Delta m_\odot^2)$		(B ₁) $= (\Delta m_\odot^2, \Delta m_{LSND}^2)$
$\circ \sin^2 2\theta_\odot \simeq 1$ $\bullet R_{atm} \simeq 0.9$ $\bullet \sin^2 2\theta_{LSND} \simeq 5 \times 10^{-5}$ $\circ r \geq 10^4$	$\circ \sin^2 2\theta_\odot \simeq 0.04$ $\bullet R_{atm} \simeq 1$ $\bullet \sin^2 2\theta_{LSND} \simeq 6 \times 10^{-5}$ $\circ r \geq 10^4$	$\bullet \sin^2 2\theta_\odot \simeq 2 \times 10^{-5}$ $\bullet R_{atm} \simeq 1$ $\circ \sin^2 2\theta_{LSND} \simeq 0.02$ $\circ r \leq 10^{-3}$
(A ₂) $(\Delta m_{32}^2, \Delta m_{21}^2) = (\Delta m_{LSND}^2, \Delta m_{atm}^2)$		(B ₂) $= (\Delta m_{atm}^2, \Delta m_{LSND}^2)$
$\circ P_{ee} \simeq 0.5$ $\circ \sin^2 2\theta_{atm} \simeq 1$ $\bullet \sin^2 2\theta_{LSND} \simeq 5 \times 10^{-5}$ $\circ r \geq 10^4$	$\bullet P_{ee} \simeq 1$ $\bullet \sin^2 2\theta_{atm} \simeq 4 \times 10^{-3}$ $\bullet \sin^2 2\theta_{LSND} \simeq 6 \times 10^{-5}$ $\circ r \geq 10^4$	$\bullet P_{ee} \simeq 1$ $\circ \sin^2 2\theta_{atm} \simeq 1$ $\circ \sin^2 2\theta_{LSND} \simeq 0.02$ $\circ r \leq 10^{-3}$

Table II. Numerical results for the special values of b_ν . The values with the underlines are input values. The values with the parentheses denote those which contradict with the observed values.

b_ν	$-\exp[i\ 1.8^\circ]$	$-(1/2)\exp[i\ 0.12^\circ]$
$r \equiv \Delta m_{32}^2/\Delta m_{21}^2$	1.950×10^4	0.03310
Δm_{32}^2	$3.51 \times 10^2\ \text{eV}^2$	<u>0.016 eV²</u>
Δm_{21}^2	<u>0.018 eV²</u>	0.483 eV ²
m_0/λ	37.5 eV	3.04 eV
m_3^ν	18.7 eV	0.707 eV
m_2^ν	0.176 eV	0.695 eV
m_1^ν	0.114 eV	0.00164 eV
$\sum m_i^\nu$	19.0 eV	1.40 eV
$\langle m_\nu \rangle$	0.00705 eV	0.00267 eV
P_{ee}	0.523	(0.990)
$\sin^2 2\theta_{atm}$	0.900	0.995
$\sin^2 2\theta_{LSND}$	(0.000048)	0.0191
$\sin^2 2\theta^{\mu\tau}$	0.211	5.4×10^{-6}
m_0	$3.12 \times 10^2\ \text{GeV}$	$3.12 \times 10^2\ \text{GeV}$
κm_0	$1.54 \times 10^{10}\ \text{GeV}$	$1.90 \times 10^{11}\ \text{GeV}$
λm_0	$2.60 \times 10^{12}\ \text{GeV}$	$3.20 \times 10^{13}\ \text{GeV}$